

AD-A043 096

MASSACHUSETTS INST OF TECH CAMBRIDGE DEPT OF PHYSICS F/G 20/13
EQUATION OF STATE AND OTHER PROPERTIES OF REALISTIC MATERIALS I--ETC(U)
DEC 76 H E STANLEY AF-AFOSR-2197-72

UNCLASSIFIED

AFOSR-TR-77-0953

NL

OF |
AD
A043096



END
DATE
FILMED
9-77
DDC

AD No. _____
DDC FILE COPY

AD A043096

(18) AFOSR-TR-77-0953

(11)

FINAL REPORT

GRANT NO. 72-2197 (15) AF-AFOSR-2197-72

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH

Title of Project

EQUATION OF STATE AND OTHER PROPERTIES OF REALISTIC MATERIALS IN THE VICINITY OF THEIR CRITICAL POINTS.

Grantee

Massachusetts Institute of Technology

Period of Award:

1 January 1972 - 31 December 1976,

(11) 31 Dec 76
(12) 28p.

DDC
RECEIVED
AUG 18 1977
C

(16) 9767

(12) 04

(10)

H. Eugene Stanley, Principal Investigator
Hermann von Helmholtz Associate Professor

Approved for public release;
distribution unlimited.

401 735

JLB

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12 (7b).
Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR- 77- 0953	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) EQUATION OF STATE AND OTHER PROPERTIES OF REALISTIC MATERIALS IN THE VICINITY OF THEIR CRITICAL POINTS		5. TYPE OF REPORT & PERIOD COVERED Scientific Final
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) H Eugene Stanley		8. CONTRACT OR GRANT NUMBER(s) AFOSR 72-2197
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physics Massachusetts Institute of Technology Cambridge MA 02139		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 9767-04 61102F
11. CONTROLLING OFFICE NAME AND ADDRESS AFOSR/NP Bolling AFB, Bldg.#410 Wash DC 20332		12. REPORT DATE
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 26
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) <p>Enginers of all sorts, from steam engines to those used in nuclear power plants, require a knowledge of a) the precise location of phase transition lines, b) "exactly" what happens when a material (such as steam) is moved across a given phase transition line. This information is essential from the engineering point of view because only with this information can one hope to have a truly efficient engine design. Phase transitions have become an extremely active field of research because of this need. Engineers, chemists, physicists, metallurgists, and mathematicians have cooperated in a multidisciplinary</p>		

DD FORM 1473

1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

effort to learn new information on this subject. These efforts can be broadly categorized as follows: a) studies focussed on critical point exponents describing the behavior of a single function as a single variable is changed; b) studies focussed on equation of state describing the behavior of a function when all thermodynamic variables are changed. The focus of this contract has been category (b), equation of state. A simple example of an equation of state is provided by a simple magnet in the vicinity of the Curie temperature T_c .

UNCLASSIFIED

CONTENTS

SIGNATURE PAGE	1
CONTENTS	2
NARRATIVE REPORT	3
INVITED TALKS BASED ON RESEARCH	
SUPPORTED BY SPONSOR	19
CONFERENCES & SUMMER SCHOOLS	
DIRECTED	20
PUBLICATIONS BASED ON RESEARCH	
SUPPORTED BY SPONSOR	
BOOKS	21
ENCYCLOPEDIA ARTICLE	21
BOOK CHAPTERS	21
JOURNAL ARTICLES	22

ACCESSION for

NTIS ☒ Write Section

ODG ☐ Buff Section

UNANNOUNCED ☐

JUSTIFICATION _____

BY _____

DISTRIBUTION/AVAILABILITY CODES

IN _____

7

FINAL REPORT

Engines of all sorts, from steam engines to those used in nuclear power plants, require a knowledge of

- a) the precise location of phase transition lines
- b) "exactly" what happens when a material (such as steam) is moved across a given phase transition line.

This information is essential from an engineering point of view because only with this information can one hope to have a truly efficient engine design.

Phase transitions have become an extremely active field of research because of this need. Engineers, chemists, physicists, metallurgists, and mathematicians have cooperated in a multidisciplinary effort to learn new information on this subject. These efforts can be broadly categorized as follows:

- a) studies focussed on critical point exponents describing the behavior of a single function as a single variable is changed.
- b) studies focussed on equation of state describing the behavior of a function when all thermodynamic variables are changed.

The focus of this contract has been category (b), equation of state. A simple example of an equation of state is provided by a simple magnet in the vicinity of the Curie temperature T_C .





FIGURE 1. PHASE TRANSITIONS & CRITICAL PHENOMENA.

- a) Phase transitions occur at all three solid lines separating different phases
- b) Critical phenomena occur at the single point (P_c, T_c) at which the first-order phase transition line separating gas and liquid phases terminates.





FIGURE 2. DIFFERENCE BETWEEN CRITICAL POINT EXPONENTS AND THE EQUATION OF STATE OF A MAGNET

- a) Phase diagram of a magnet, showing information learned by critical point exponents along 2 paths
- b) By contrast, the equation of state provides information in the entire phase diagram near the critical point ($H=0, T=T_c$).

The phase diagram of the simple magnet is provided in Fig.2. There is a line of first order transitions which terminates in a critical point at $T=T_c$, $H=0$ where H is the applied magnetic field. Studies of category (a) concern the determination of critical exponents along paths (i) and (ii) of Fig.2a. For example, if the function in question is the spontaneous magnetization, $M = M(H,T)$, then along path (i)

$$M = M(H=0,T) \sim t^\beta \quad (1a)$$

while along path (ii),

$$M = M(H,T=T_c) \sim H^{1/\delta} \quad (1b)$$

thereby defining the two critical exponents β and δ . Here we have used the notation $t \equiv (T-T_c)/T_c$ to denote the reduced temperature (i.e., $t=0$ at the critical point).

Studies of category (b) concern the determination of the function in the entire region near the critical point. For our example of the magnetization function $M(H,T)$, this means that we must determine M everywhere in the region shown shaded in Fig.2b.

Clearly studies of category (b) provide more information than studies of category (a), and therefore it is not surprising that they are more difficult to carry out.

The principal difficulty stems from the fact that very close to the critical point the material undergoes fluctuations that are not only characterized by their size but also are characterized by the fact that the fluctuations

of all wavelengths are simultaneously present. This means that any approximation procedure of the sort familiar from solid state physics is doomed to failure.

Our research group has participated in the development of an approach to the development of a theory of the equation of state. This approach permits the accurate calculation of the equation of state for a wide range of materials.

Specifically, it has^{been} found that very near the critical point, the equation of state of a wide range of materials is a "generalized homogeneous function" (GHF). A function of two variables H and t is a GHF if there exist two numbers (called scaling powers) a_H and a_t such that for all positive values of some arbitrary parameter L ,

$$f(L^{a_H} H, L^{a_t} t) = L f(H, t) \quad (2)$$

Specifically, the function $f(H, t)$ is the Gibbs potential. Since every thermodynamic function is related by a series of Legendre transformations and partial differentiations to the Gibbs potential, Equation (2) has far-reaching implications. Specifically, the magnetization function $M(H, t)$ is related to the Gibbs potential by the relation

$$M = \left[\partial f(H, t) / \partial H \right] \quad \text{at } t = \text{const} \quad (3)$$

Therefore from (2) and (3) it follows that $M(H, t)$ is also a GHF; that is,

$$M(L^{a_H} H, L^{a_t} t) = L^{1-a_H} M(H, t) \quad (4)$$

The physical content of Equation (4) is that the equation of state $M(H,t)$ is constrained rather severely. To see this, consider the equation of state for a typical material. This consists of a series of values of $M(T)$ for various values of H . That is, for each value of H , one must measure $M(T)$. Such data are shown schematically in Fig.3, as a family of curves, one for each value of H .

Now Equation (4) implies that the entire family of curves collapses onto one single curve providing the experimental data are plotted in a "scaled form". To see this, note from (4) that L is an arbitrary number, and therefore can be chosen as

$$L = (1/H)^{1/a_H} \quad (5)$$

Substituting (5) into (4), we obtain

$$\frac{M(H,t)}{H^{(1-a_H)/a_H}} = M(1, t / H^{a_t/a_H}) \quad (6)$$

But Equation (6) states that if one plots "scaled magnetization"

$$\tilde{M} \equiv M / H^{(1-a_H)/a_H}$$

as a function of "scaled temperature"

$$\tilde{t} \equiv t / H^{a_t/a_H} \quad (7)$$

then one finds that the entire family of curves in Fig.3a "collapse" onto a single curve in Fig.3b.

This single curve is called the "scaled equation of state" or the "scaling function" for short. The experimental verification of the scaled equation of state is rather




FIGURE 3.

Illustration of "data collapsing" onto a single "scaled equation of state".




FIGURE 4.

Data collapsing for 5 different materials onto a single scaling function. Solid curve = calculations.

striking, as shown in Figure 4. Here we have plotted experimental data taken on five widely diverse materials. The first material, CrBr_3 , has considerable lattice anisotropy (the coupling strengths in the z lattice direction are about 17 times weaker than those along the x or y directions). The second material, EuO , has significant second-neighbor interactions, and these are probably of the opposite sign to the nearest-neighbor interactions. The third material, Ni , is probably an itinerant electron ferromagnet, while the fourth material Pd_3Fe is an alloy and the fifth material YIG is actually a ferrimagnet. All five materials display data collapsing, thereby providing experimental confirmation of the scaling equation of state.

From Figure 4, we see that all these data fall on one and only one "scaling function"--i.e., all five materials have the identical scaled equation of state! This fact suggests that while the equation of state is strongly dependent on the details of intermolecular interactions for most regions of the phase diagram, at least near the critical point it appears to be independent of such details for otherwise five such widely diverse materials could hardly obey the identical scaled equation of state!

Such apparent universality has provided impetus for a variety of calculations during the tenure of this grant. Specifically, we have developed the method of high-temperature

series expansions in such a fashion that it can be used to provide information on the equation of state of a wide variety of materials. Then we have proceeded to calculate the equation of state and to compare the calculated scaling functions with experimental data. The calculated function is shown in Figure 4 as the solid curve, and we note that the agreement is rather satisfactory--especially considering the fact that there are no free parameters in the theory.

It is perhaps not entirely inappropriate to record the fact that this work has led to invitations to address three major international meetings:

- a) the International Conference on Low-Temperature Physics
- b) the International Congress on Magnetism, Moscow, USSR
- c) the International Enrico Fermi School of Physics, Varenna, Italy

So much for studies of ordinary critical points. It is becoming increasingly appreciated that more complex materials display much more complex critical points. For example, Soviet work reported first in the Russian Journal of Physical Chemistry has stimulated a tremendous increase in our understanding of such complex materials. Specifically, the Soviet work considered materials that consisted of three-component fluids mixed in such proportions that a "tricritical point" can occur.

Subsequent to the early work by Soviet authors, tricritical points have been detected experimentally in an extremely wide range of materials, including not only multicomponent fluid mixtures but also magnetic materials, superconductors, liquid crystals, and so forth.

Our work began with the study of metamagnetic materials, which are characterized by a sudden discontinuous change in magnetization as a function of applied magnetic field. These materials are among the simplest from a theoretical point of view. In particular, colleagues from the Division of Engineering & Applied Physics of Yale University (W.P.Wolf and co-workers) have made detailed measurements of the equations of state of dysprosium aluminum garnet (DAG), one such metamagnet. A natural theoretical question is the following: "can these data be interpreted in terms of a unifying scaling hypothesis?" To this time, there had been no experimental tests of the scaling hypothesis at tricritical points. However in light of the success of the scaled equation of state at an ordinary critical point, we were optimistic and decided to analyze the Yale data on DAG.

We found that the data did indeed collapse onto a single curve, but the properties of this curve were more like those of an ordinary critical point than a tricritical point. Thus our results remained a mystery until Blume and co-workers at Brookhaven National Laboratory realized that

the experimental data were obtained under conditions in which the applied field coupled to the order parameter and thereby moved one off the H-T plane. The experiments have recently been repeated with a different orientation of the H field, and the predictions of tricritical scaling confirmed.

The experimental data are plotted in Fig.5, and the general phase diagram of a tricritical point is shown in Fig.6.

In addition to "ordinary" critical points and "tricritical points", we have considered the possibility of even more complex critical points. For example, a tricritical point occurs at the intersection of lines of ordinary critical points. This leads to the question "What sort of critical point would occur at the intersection of lines of tricritical points?"

To answer this question, we have developed a classification of complex critical points that is fairly general (though probably not sufficiently general to handle the most complex materials in nature). Specifically, we have introduced the concept of the "order" of a critical point. Ordinary critical points are of order 2, while tricritical points are of order 3 and the point of intersection of critical lines is of order 4. A metamagnet in the presence of both a direct and a staggered magnetic field can be shown to exhibit a critical point of order 4, and we made calculations of critical exponents for such a system.




FIGURE 5.

Experimental data of the Yale group on the metamagnetic material dysprosium aluminum garnet (DAG). The data collapse onto a single curve, the scaled equation of state. The data analysis was carried out under the auspices of this grant.




FIGURE 6.

Phase diagram for a typical model metamagnet. Note that three different second order lines of "ordinary" critical points intersect at a "tricritical point" (TCP).

A final class of "realistic materials" will be discussed. These are materials which in one sense or another can be considered to have an effective dimensionality less than 3. Such materials are likely to become extremely important in engineering applications. For example, graphite is in many senses effectively 2-dimensional, and this near 2-dimensionality results in many of its unusual mechanical properties. Similarly, the search for high-temperature superconductors has focussed in part upon materials that are quasi-2-dimensional.

Therefore we have focussed our attention upon a class of materials in which the interactions between spins in one lattice direction are R times those in the remaining 2 directions. For R very small, the system is quasi-2-dimensional, while for R very large the system is quasi-1-dimensional.

Experimental work on materials with R very small has suggested the possible occurrence of a "Stanley-Kaplan phase transition" to a novel low-temperature phase in which there is an infinite susceptibility but no long-range order.

We have developed a set of rigorous relations that relate the measured properties of the full three-dimensional system (i.e., $R \neq 0$) to the two-dimensional system ($R=0$). Similarly, we have been able to predict the location of the crossover from effectively two-dimensional behavior far above the critical temperature to a limiting form of

three-dimensional behavior very close to the critical temperature. This work has led to a joint collaboration with L.J.deJongh at the Kamerlingh Onnes Laboratorium, Leiden. deJongh carried out experiments that verified the predictions of the crossover theory, and the results were published jointly with the Principal Investigator in a recent issue of Physical Review Letters.

In summary, then, we have studied a wide variety of materials that are relatively complex compared with the idealized systems that are usually "first" studied by theorists. These systems have included materials such as DAG that display higher order critical points, and materials that display quasi-1-dimensional or quasi-2-dimensional order. It has been found that these materials have an equation of state that obeys a remarkably simple form when expressed in terms of the proper variables.




FIGURE 7.

Analysis of crossover phenomena in quasi-2-dimensional magnetic materials. A system changes its critical properties as the critical point is approached arbitrarily closely, in accordance with certain rigorous results of Liu and Stanley (1973) and confirmed experimentally by deJongh and Stanley (1976).

INVITED TALKS BASED ON RESEARCH SUPPORTED BY THE SPONSOR:

1. Invited speaker (with S. Milosevic, D. Karo, and R. Krasnow), Thirteenth International Conference on Low-Temperature Physics ("LT-13"). Boulder, Colorado, August 1972.
2. Principal Invited Speaker (6 intro. lectures), Enrico Fermi Summer School on Local Phenomena near Phase Transitions, Varenna, Italy, July 1973.
3. Invited speaker (with L. Liu), International Workshop on Physics in Less than Three Dimensions, July 1973.
4. Invited speaker, International workshop on Applications of Statistical Mechanics to Cooperative Phenomena, Dartmouth College, August 1973.
5. Invited speaker (with S. Milosevic), International Congress on Magnetism, Moscow, U.S.S.R., August 1973.
6. Invited speaker, Conference on Cooperative Phenomena, Portoroz, Yugoslavia, June 1974.
7. Invited speaker, Latin American School of Physics, Mexico City, July 1974.
8. Invited speaker, AMERICAN PHYSICAL SOCIETY, May 1975 (Kent, Ohio).
9. Invited speaker, NOZAWA SUMMER SCHOOL IN SOLID STATE PHYSICS, NOzawa, Nagano Pref., Japan, July 1975.
10. Invited speaker, INTL. CONF. ON LOW-LYING LATTICE VIBRATIONAL MODES AND THEIR RELATIONSHIP TO SUPER-CONDUCTIVITY AND FERROELECTRICITY, San Juan, P.R., 1-5 Dec. 1975.

CONFERENCES AND SUMMER SCHOOLS DIRECTED:

1. International Organizing Committee (with P.W. Anderson, R. Blinc, H.Z. Cummins, J.A. Krumhansl, B.T. Matthias, J.R. Schrieffer, H.G. Smith, and G. Shirane), INTERNATIONAL CONFERENCE ON LOW-LYING LATTICE VIBRATIONAL MODES AND THEIR RELATIONSHIP TO SUPER-CONDUCTIVITY AND FERROELECTRICITY, 1-5 December 1975. San Juam, P.R.
2. Director, Summer Program on Phase Transitions and Critical Phenomena, M.I.T., Cambridge, MA, July 1970, June 1971, June, 1972, June 1973, June 1974 and June 1975.

PUBLICATIONS BASED ON RESEARCH SUPPORTED BY THE SPONSOR:

BOOKS:

1. H.E. Stanley, Introduction to Phase Transitions and Critical Phenomena, a book in the "International Series of Monographs on Physics" of Oxford University Press, Oxford and New York 1971 (308) pages.

Translations:

Russian:

Translated by S.V. Vonsovsky
(MIR, Moscow, 1973)

Japanese:

(Soteni to Rinkai
Gensho) Translated by K. Matsuno
(Tokyo-Tosho, Tokyo, 1974)

2. H.E. Stanley, Ed., Cooperative Phenomena near Phase Transitions, M.I.T. Press, Cambridge, 1973.
Paper and Hardcover editions, 308 pages.

ENCYCLOPEDIA ARTICLE

3. H.E. Stanley, "Critical Phenomena", in Encyclopedia of Physics (Ed. R.M. Besancon). Van Nostrand and Reinhold Publ. Co., N.Y., 1974.

BOOK CHAPTERS

4. H.E. Stanley, "Scaling Laws and Universality -- or Statistical Mechanics is not Dead!" In Statistical Mechanics and Field Theory, (Ed. R. Sen), Keter Publ. Co., Jerusalem, 1972, p. 225-267.
5. H.E. Stanley, "The D-Vector Model of 'Universality Hamiltonian': Properties of Isotropically-interacting D-dimensional Classical Spins". In Phase Transitions and Critical Phenomena (Eds. C. Domb and M.S. Green) Academic Press, London, 1974. Vol. III, Chapter 7, pages 485-567.

6. H.E. Stanley, T.S. Chang, F. Harbus, and L.L. Liu, "Five Introductory Lectures on Critical Phenomena in Simple and Complex Systems: The Unifying Hypotheses of Scaling and Universality", In Proc. 1973 Enrico Fermi Varenna School: Course 59-Local Properties at Phase Transitions, K.A. Müller, and A. Rigamonti, Ed. (North Holland, Amsterdam, 1975) pp. 45-137.

JOURNAL ARTICLES

7. L.L. Liu and H.E. Stanley, "Some Results Concerning the Crossover Behavior of Quasi-two-dimensional and Quasi-one-dimensional Systems", Physical Review Letters 29, 927-931 (1972).
8. L.L. Liu and H.E. Stanley, "Quasi-one Dimensional and Quasi-two-Dimensional Magnetic Systems: Determination of Crossover Temperature and Scaling with Anisotropy Parameters", Physical Review B 8, 2279-2298 (1973).
9. L.J. deJongh and H.E. Stanley, "Lattice-Dimensionality Crossover Effects in Quasi-d-Dimensional Magnetic Materials", Physical Review Letters 36, 817-820 (April 1976).
10. D.N. Lambeth and H.E. Stanley, "Models for Quasi-Two-Dimensional Helium and Magnets", Physical Review B 12, 5302-5314 (1 December 1975).
11. A. Hankey and H.E. Stanley, "Systematic Application of Generalized Homogeneous Functions to Static Scaling, Dynamic Scaling, and Universality", Physical Review B 6, 3515-3542 (1972).
12. J. Nicoll, T.S. Chang, H.E. Stanley, and A. Hankey, "Scaling Laws for Fluid Systems Using Generalized Homogeneous Functions of Strong and Weak Variables", Physical Review B 11, 1176-1187 (1975).
13. S. Milosevic and H.E. Stanley, "Calculation of the Scaling Function for the Heisenberg Model", Physical Review B 5, 2526-2529 (1972).

14. S. Milosevic and H.E. Stanley, "Equation of State Near the Critical Point I. Calculation of the Scaling Function for the $S=1/2$ and $S=\infty$ Heisenberg Models Using High-Temperature Expansions", Physical Review B6, 986-1001 (1972).
15. S. Milosevic and H.E. Stanley, "Equation of State Near the Critical Point. II. Comparison with Experiment and Possible Universality with Respect to Lattice Structure and Spin Quantum Number", Physical Review B6, 1002-1008 (1972).
16. S. Milosevic, D. Karo, R. Krasnow, and H.E. Stanley, "Magnetic Equations of State near the Critical Point", Invited talk, Proc. Thirteenth International Low-Temperature, E. Hammel, W. O'Sullivan, and K. Timmerhaus, Eds. (Plenum Press, N.Y., 1974, Vol. 2, pp. 295-321).
17. S. Milosevic, D. Karo, R. Krasnow, and H.E. Stanley "Scaling Functions: Comparison of Theory and Experiment", Invited talk, Proc. International Congress on Magnetism, Moscow, U.S.S.R., 1973. Vol. I (1). pp 105-116.
18. S. Milosevic, and H.E. Stanley, "Scaling Functions of Magnets". In Proc. 1973 Enrico Fermi Summer School: Course 59--Local Properties at Phase Transitions, K.A. Muller and A. Rigamonti, eds. (North Holland, Amsterdam, 1975). pp 773-785.
19. R. Krasnow and H.E. Stanley, "Specific Heat Scaling Functions for Ising and Heisenberg Models and Comparison with Nickel Data" Physical Review B8, 332-338 (1973)
20. F. Harbus and H.E. Stanley, "General - R High-Temperature Series for the Susceptibility, Second Moment, and specific Heat of the sc and fcc Ising Models with Lattice Anisotropy", Physics Review B7, 365-370, (1973).
21. R. Krasnow, F. Harbus, L.L. Liu, and H.E. Stanley, "Scaling with Respect to a parameter for the Gibbs Potential and Pair Correlation Function of the $S=1/2$ Ising Model with Directional Anisotropy", Physics Review B7, 370-379 (1973).
22. G. Paul and H.E. Stanley, "A Partial Test of the Universality Hypothesis. The Case of Next-Nearest Neighbor Interactions", Physical Review B5, 3715-3726 (1972).

23. G. Paul and H.E. Stanley, "A Partial Test of the Universality Hypothesis. The Case of Next-Nearest Neighbor Interactions", Physical Review B5, 3715-3726 (1972).
24. D. Lambeth, M.H. Lee and H.E. Stanley, "High-Temperature Series for the B-Site Spinel and Diamond Lattices and the Universality Question," Journal of Chemical Physics 60, 770-780 (1974).
25. T.S. Chang, A. Hankey and H.E. Stanley, "Proposal for Classification of Critical Points by Order", Physics Letters 44A, 25-26 (1973).
26. A. Hankey, T.S. Chang and H.E. Stanley, "Two Soluble Models of Magnetic Systems Showing Critical Points of Higher Order", Proceedings of the 1972 Conference on Magnetism and Magnetic Materials, A.I.P. Conference Publication No. 10, p. 889-892 (1973).
27. F. Harbus, A. Hankey, H.E. Stanley, and T.S. Chang, "System Exhibiting a Critical Point of Order Four: Ising Planes with Variable Interplanar Interactions", Physical Review B8, 2273-2278 (1973).
28. F. Harbus and H.E. Stanley, "Ising Model Metamagnet and Tricritical Susceptibility Exponent", Physical Review Letters 29, 58-62 (1972).
29. F. Harbus and H.E. Stanley, "A High-Temperature Series Study of Models Displaying Tricritical Behavior. I. Ferromagnetic Planes Coupled Antiferromagnetically", Physical Review B8, 1141-1156 (1973).
30. F. Harbus and H.E. Stanley, "A High-Temperature Series Study of Models Displaying Tricritical Behavior. II. A Nearest-Neighbor Ising Antiferromagnet with Next-Nearest-Neighbor Ferromagnetic Interactions", Physical Review B8, 1156-1167 (1973).

31. M. Wortis, F. Harbus, and H.E. Stanley,
"Tricritical Behavior of the Ising Anti-ferromagnet with Next-Nearest-Neighbor Ferromagnetic Interactions: Mean-Field-Like Tricritical Exponents?", Physical Review B11, 2689-2692 (April 1975).
32. A. Hankey, H.E. Stanley, and T.S. Chang,
"Geometric Predictions of Scaling at Tricritical points", Physical Review Letters 29, 278-281 (1972).
33. T.S. Chang, A. Hankey, and H.E. Stanley,
"Double-Power Scaling Functions near Tricritical Points", Physical Review B7, 4263-4266 (1973).
34. G.F. Tuthill, F. Harbus, H.E. Stanley,
"Test of Tricritical Point Scaling in DAG"
Physical Review Letters 31, 527-530 (1973).
35. F. Harbus and H.E. Stanley, "Use of Scaling Theory to Predict Amplitudes: Verification of Double Power Law Behavior for Crossover of Lattice Dimensionality", Physical Review B8, 2268-2272 (1973).
36. T.S. Chang, A. Hankey, and H.E. Stanley,
"Generalized Scaling Hypothesis in Multicomponent Systems. I. Classification of Critical Points by Order and Scaling at Tricritical Points", Physics Review B8, 346-364 (1973).
37. A. Hankey, T.S. Chang and H.E. Stanley,
"Generalized Scaling Hypothesis in Multicomponent Systems. II. The Scaling Hypothesis at CRitical PRints of Arbitrary Order", Physical Review B8, 1178-1184, (1973).
38. A. Hankey, T.S. Chang, and H.E. Stanley,
"Tricritical Points in Multi-component Fluid Mixtures: A Geometric Interpretation", Physical Review A9, 2573-2578 (1974).
39. L.L. Liu and H.E. Stanley, "Exact Solution in an external Magnetic Field of Ising Models with Three-Spin Interactions", Physical Review B10, 2958-2961 (1 October 1974).

40. L.L. Liu, R.I. Joseph, and H.E. Stanley, "New Inequalities among the Critical-Point Exponents for the Spin-Spin and Energy-Energy Correlation Functions", Physical Review B6, 1963-1968 (1972).
41. L.L. Liu and H.E. Stanley, "Divergence of Correlation Length along the Critical Isotherm". Physical Review B7, 3241-3244(1973).
42. P. Horn, R.D. Parks, D. Lambeth, and H.E. Stanley, "Critical Magnetic Susceptibility in the presence of Long Range Forces," Physical Review B9, 316-322 (1974).
43. H. Birecki, H.E. Stanley, and Y. Shapira, "Method of Calculating High-Temperature Series Expansions Exact in the External Magnetic Field; Application to the Two-Spin Correlation Function and Magnetization of the Classical Heisenberg Magnet", Physical Review B8, 2327-2335 (1973).
44. H.E. Stanley, R.J. Birgeneau, P.J. Reynolds, and J.F. Nwcoll, "Thermally Driven Phase Transitions near the Percolation Threshold in Two Dimensions", J. Phys. C: Solid State Physics, 9, L553-560 (28 October 1976).
45. P.J. Reynolds, W. Klein, and H.E. Stanley, "Renormalization Group for Site and Bond Percolation", Journal of Physics C: Solid State Physics 10, L167-L 172 (28 April 1977).
46. H.E. Stanley, "Cluster Shapes at the Percolation Threshold: Connection Between Critical Point Exponents and the Fractal Dimensionality of Clusters" Journal of Physics A (submitted for publication).